

Computer Graphics

Lecture 14

2D Transformation Matrices

Homogeneous Coordinates



Computer Graphics

Lecture 14


2D Transformation Matrices

Homogeneous Coordinates

Announcements

- Final projects will be done in groups of
 - 480: 2 or 3
 - 580: 2

Situation: Bunny is sad.



Bunny is sad because it can't move.

Today: Make bunny happy

- How can we manipulate objects in the scene to
 - put them in the right position?
 - scale them to the right size?
 - orient them in the right direction?

Our answer: matrices.



Geometric Transformations

- To the notes!

Linear transformation gallery

- Uniform scale $\begin{bmatrix} s & 0 \\ 0 & s \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} sx \\ sy \end{bmatrix}$



$$\begin{bmatrix} 1.5 & 0 \\ 0 & 1.5 \end{bmatrix}$$



Linear transformation gallery

- Shear $\begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + ay \\ y \end{bmatrix}$

$$\begin{bmatrix} 1 & 0.5 \\ 0 & 1 \end{bmatrix}$$





Linear transformation gallery

- Nonuniform scale

$$\begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} s_x x \\ s_y y \end{bmatrix}$$


$$\begin{bmatrix} 1.5 & 0 \\ 0 & 0.8 \end{bmatrix}$$



Linear transformation gallery

- Reflection
 - can consider it a special case of nonuniform scale



$$\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$



Linear transformation gallery

- Rotation
$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \cos \theta - y \sin \theta \\ x \sin \theta + y \cos \theta \end{bmatrix}$$

$$\begin{bmatrix} 0.866 & -0.5 \\ 0.5 & 0.866 \end{bmatrix}$$



2D Matrix Transformations: Properties

- linear
- closed under composition
- associative
- not commutative
- applied right-to-left

Composing transformations

- Want to move an object, then move it some more
 - $\mathbf{p} \rightarrow T(\mathbf{p}) \rightarrow S(T(\mathbf{p})) = (S \circ T)(\mathbf{p})$
- We need to represent $S \circ T$ (“S compose T”)
 - and would like to use the same representation as for S and T
- Translation easy
 - $T(\mathbf{p}) = \mathbf{p} + \mathbf{u}_T; S(\mathbf{p}) = \mathbf{p} + \mathbf{u}_S$
 $(S \circ T)(\mathbf{p}) = \mathbf{p} + (\mathbf{u}_T + \mathbf{u}_S)$
- Translation by \mathbf{u}_T then by \mathbf{u}_S is translation by $\mathbf{u}_T + \mathbf{u}_S$
 - commutative!

Composing transformations

- Linear transformations also straightforward

- $T(\mathbf{p}) = M_T \mathbf{p}; S(\mathbf{p}) = M_S \mathbf{p}$

- $(S \circ T)(\mathbf{p}) = M_S M_T \mathbf{p}$

- Transforming first by M_T then by M_S is the same as transforming by $M_S M_T$

- only sometimes commutative

- e.g. rotations & uniform scales
 - e.g. non-uniform scales w/o rotation

- Note $M_S M_T$, or $S \circ T$, is T first, then S

Combining linear with translation

- Need to use both in single framework
- Can represent arbitrary seq. as $T(\mathbf{p}) = M\mathbf{p} + \mathbf{u}$
 - $T(\mathbf{p}) = M_T\mathbf{p} + \mathbf{u}_T$
 - $S(\mathbf{p}) = M_S\mathbf{p} + \mathbf{u}_S$
 - $(S \circ T)(\mathbf{p}) = M_S(M_T\mathbf{p} + \mathbf{u}_T) + \mathbf{u}_S$
 $= (M_S M_T)\mathbf{p} + (M_S\mathbf{u}_T + \mathbf{u}_S)$
 - e.g. $S(T(\mathbf{0})) = S(\mathbf{u}_T)$
- Transforming by M_T and \mathbf{u}_T , then by M_S and \mathbf{u}_S , is the same as transforming by $M_S M_T$ and $\mathbf{u}_S + M_S\mathbf{u}_T$
 - This will work but is a little awkward

Homogeneous coordinates

- A trick for representing the foregoing more elegantly
- Extra component w for vectors, extra row/column for matrices
 - for affine, can always keep $w = 1$
- Represent linear transformations with dummy extra row and column

$$\begin{bmatrix} a & b & 0 \\ c & d & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} ax + by \\ cx + dy \\ 1 \end{bmatrix}$$

Homogeneous coordinates

- Represent translation using the extra column

$$\begin{bmatrix} 1 & 0 & t \\ 0 & 1 & s \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x + t \\ y + s \\ 1 \end{bmatrix}$$

Homogeneous coordinates

- Composition just works, by 3x3 matrix multiplication

$$\begin{bmatrix} M_S & \mathbf{u}_S \\ 0 & 1 \end{bmatrix} \begin{bmatrix} M_T & \mathbf{u}_T \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \mathbf{p} \\ 1 \end{bmatrix} \\ = \begin{bmatrix} (M_S M_T) \mathbf{p} + (M_S \mathbf{u}_T + \mathbf{u}_S) \\ 1 \end{bmatrix}$$

- This is exactly the same as carrying around M and \mathbf{u}
 - but cleaner
 - and generalizes in useful ways as we'll see later

Affine transformations

- The set of transformations we have been looking at is known as the “affine” transformations
 - straight lines preserved; parallel lines preserved
 - ratios of lengths along lines preserved (midpoints preserved)

