


CSCI 497P/597P: Computer Vision

Scott Wehrwein

Cameras and Panoramas



Reading


- Szeliski: Chapter 2.1.5-6, 6.3

Goals

- Know the properties and projection matrix associated with **orthographic projection**.
- Understand the concept of **focal length** and its relationship with the projection matrix.
- Understand the causes and corrections for perspective and radial lens distortion.
- Understand how to construct a planar panorama by warping images to a common projection plane using homographies.


Announcements

Pinhole camera




- Add a barrier to block off most of the rays
 - This reduces blurring
 - The opening known as the **aperture**
 - How does the image relate to the scene?

Modeling projection



- The coordinate system
 - We will use the pinhole model as an approximation
 - Put the optical center (**C**enter **O**f **P**rojection) at the origin
 - Put the image plane (**P**rojection **P**lane) *in front* of the COP
 - Why?
 - The camera looks down the *negative* z axis
 - we like this if we want right-handed-coordinates

Modeling projection



- Projection equations

- Compute intersection with PP of ray from (x,y,z) to COP
- Derived using similar triangles (on board)

$$(x, y, z) \rightarrow \left(-d\frac{x}{z}, -d\frac{y}{z}, -d\right)$$

- We get the projection by throwing out the last coordinate:

$$(x, y, z) \rightarrow \left(-d\frac{x}{z}, -d\frac{y}{z}\right)$$

Modeling projection

- Is this a linear transformation?
 - no—division by z is nonlinear

Homogeneous coordinates to the rescue—again!

$$(x, y) \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

homogeneous image
coordinates

$$(x, y, z) \Rightarrow \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$

homogeneous scene
coordinates

Converting *from* homogeneous coordinates

$$\begin{bmatrix} x \\ y \\ w \end{bmatrix} \Rightarrow (x/w, y/w)$$

$$\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \Rightarrow (x/w, y/w, z/w)$$

Perspective Projection

Projection is a matrix multiply using homogeneous coordinates:

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1/d & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} x \\ y \\ -z/d \end{bmatrix} \Rightarrow \left(-d \frac{x}{z}, -d \frac{y}{z} \right)$$

divide by third coordinate

This is known as **perspective projection**

- The matrix is the **projection matrix**
- (Can also represent as a 4x4 matrix – OpenGL does something like this)

Perspective Projection

- How does scaling the projection matrix change the transformation?

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1/d & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} x \\ y \\ -z/d \end{bmatrix} \Rightarrow \left(-d\frac{x}{z}, -d\frac{y}{z}\right)$$

$$\begin{bmatrix} -d & 0 & 0 & 0 \\ 0 & -d & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} -dx \\ -dy \\ z \end{bmatrix} \Rightarrow \left(-d\frac{x}{z}, -d\frac{y}{z}\right)$$

Perspective Projection

$$\begin{bmatrix} -d & 0 & 0 & 0 \\ 0 & -d & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} -dx \\ -dy \\ z \end{bmatrix} \Rightarrow \left(-d\frac{x}{z}, -d\frac{y}{z}\right)$$

What is the meaning of d?

d = Focal length

- Can think of as “zoom”



24mm



50mm



200mm




800mm




- Also related to *field of view*


Field of view

APS-C Crop Body Measurement Table

Lens	After 1.62 Multiplier	APS-C Sensor (1.62 lens multiplier) Canon 60D, 7D, 70D, T3i, T4i	Hand Positions
18mm	29.16mm	Three hands wide at full arms length.	
28mm	45.36mm	Slightly less than two hands wide at full arms length.	
35mm	56.7mm	One hand + width of one fist at full arms length.	
50mm	81mm	One hand wide + width of thumb at full arms length.	
55mm	89.1mm	Slightly less than one hand wide at full arms length.	
85mm	137.7mm	Inside edge of thumb to tip of forefinger wide with hand in "L" shape, thumb up.	



Focal length in practice



24mm




50mm



135mm



Focal length = cropping



24mm



50mm




135mm



Focal length vs. viewpoint

- Telephoto makes it easier to select background (a small change in viewpoint is a big change in background).



Grand-angle 24 mm



Normal 50 mm



Longue focale 135 mm



Fredo Durand




Wide angle



Standard



Telephoto



<http://petapixel.com/2013/01/11/how-focal-length-affects-your-subjects-apparent-weight-as-seen-with-a-cat/>

Fredo Durand

Perspective Projection


$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1/d & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} x \\ y \\ -z/d \end{bmatrix} \Rightarrow \left(-d\frac{x}{z}, -d\frac{y}{z}\right)$$

divide by third coordinate

What does it mean to set $f(d)$ to infinity?


Orthographic projection

- Special case of perspective projection
 - Distance from the COP to the PP is infinite
 - In other words, $f = \infty$




$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \Rightarrow (x, y)$$

Orthographic projection



Perspective projection




Perspective distortion

- Problem for architectural photography: converging verticals




Perspective distortion


- Problem for architectural photography: converging verticals



Tilting the camera upwards results in converging verticals





Keeping the camera level, with an ordinary lens, captures only the bottom portion of the building




Shifting the lens upwards results in a picture of the entire subject

- Solution: view camera (lens shifted w.r.t. film)



Perspective distortion


- Problem for architectural photography: converging verticals
- Result:



Perspective distortion: People



Lens Distortion



- Radial distortion of the image
 - Caused by imperfect lenses
 - Deviations are most noticeable for rays that pass through the edge of the lens



Modeling distortion

Project $(\hat{x}, \hat{y}, \hat{z})$
to “normalized”
image coordinates

$$x'_n = \hat{x} / \hat{z}$$
$$y'_n = \hat{y} / \hat{z}$$

Apply radial distortion

$$r^2 = x_n'^2 + y_n'^2$$
$$x'_d = x'_n(1 + \kappa_1 r^2 + \kappa_2 r^4)$$
$$y'_d = y'_n(1 + \kappa_1 r^2 + \kappa_2 r^4)$$


Apply focal length
translate image center

$$x' = f x'_d + x_c$$
$$y' = f y'_d + y_c$$

- To model lens distortion
 - Use above projection operation instead of standard projection matrix multiplication


Dimensionality Reduction Machine (3D to 2D)

3D world



Point of observation

2D image



What have we lost?



- Angles
- Distances (lengths)

Projection properties

- Many-to-one: any points along same ray map to same point in image
- Points $\rightarrow$ points
- Lines $\rightarrow$ lines (collinearity is preserved)
 - But line through focal point projects to a point
- Planes $\rightarrow$ planes (or half-planes)
 - But plane through focal point projects to line


Projection properties

- Parallel lines converge at a vanishing point
 - Each direction in space has its own vanishing point
 - But parallels parallel to the image plane remain parallel





Questions?

Back to panoramas



Can we use homographies to create a 360 panorama?

Homographies project images onto a common plane.



each image is warped
with a homography $\mathbf{H}$


We'll see what this homography means in
terms of cameras and projection later.

First -- Can't create a 360 panorama this way...

mosaic **Projection Plane**

Panoramas

- What if you want a 360° field of view?




mosaic **Projection Sphere**



Unwrapping a sphere


Credit: JHT's Planetary Pixel Emporium



Spherical reprojection




input



$f = 200$ (pixels)



$f = 400$



$f = 800$

- Map image to spherical coordinates
 - need to know the focal length